

Quantum dots: transport

- [1] G. Kießlich, A. Wacker, and E. Schöll: *Theory of nonlinear transport for ensembles of quantum dots*, in *Semiconductor Nanostructures*, edited by D. Bimberg (Springer, Berlin, 2008), pp. 211–220.
- [2] G. Kießlich, E. Schöll, F. Hohls, and R. J. Haug: *Coulomb-mediated electron bunching in tunneling through coupled quantum dots*, *Phys. stat. sol. (c)* **5**, 166 (2008).
- [3] G. Kießlich, E. Schöll, T. Brandes, F. Hohls, and R. J. Haug: *Noise enhancement due to quantum coherence in coupled quantum dots*, *Phys. Rev. Lett.* **99**, 206602 (2007).
- [4] C. Gnodtke, G. Kießlich, A. Wacker, and E. Schöll: *Phonon-assisted tunneling through quantum dot stacks*, *Phys. Rev. B* **73**, 115338 (2006).
- [5] G. Kießlich, P. Samuelsson, A. Wacker, and E. Schöll: *Counting statistics and decoherence in coupled quantum dots*, *Phys. Rev. B* **73**, 033312 (2006).
- [6] H. Sprekeler, G. Kießlich, A. Wacker, and E. Schöll: *Coulomb effects in tunneling through a quantum dot stack*, *Phys. Rev. B* **69**, 125328 (2004).
- [7] G. Kießlich, H. Sprekeler, A. Wacker, and E. Schöll: *Positive correlations in tunnelling through coupled quantum dots*, *Semicond. Sci. Technol.* **19**, S37 (2004).
- [8] R. Wetzler, R. Kunert, A. Wacker, and E. Schöll: *Inhomogeneous charging and screening effects in semiconductor quantum dot arrays*, *New J. Phys.* **6**, 81 (2004).
- [9] R. Wetzler, A. Wacker, and E. Schöll: *Non-local Auger effect in quantum dot devices*, *Semicond. Sci. Technol.* **19**, S43 (2004).
- [10] R. Wetzler, A. Wacker, and E. Schöll: *Coulomb scattering with remote continuum states in quantum dot devices*, *J. Appl. Phys.* **95**, 7966 (2004).
- [11] R. Wetzler, A. Wacker, and E. Schöll: *Self-consistent Coulomb effects and charge distribution of quantum dot arrays*, *Phys. Rev. B* **68**, 045323 (2003).
- [12] G. Kießlich, A. Wacker, and E. Schöll: *Shot noise of coupled semiconductor quantum dots*, *Phys. Rev. B* **68**, 125320 (2003).
- [13] G. Kießlich, A. Wacker, E. Schöll, A. Nauen, F. Hohls, and R. J. Haug: *Shot noise in tunneling through a quantum dot array*, *phys. status solidi (c)* **0**, 1293 (2003).
- [14] G. Kießlich, A. Wacker, E. Schöll, S. A. Vitusevich, A. E. Belyaev, S. V. Danylyuk, A. Förster, N. Klein, and M. Henini: *Nonlinear charging effect of quantum dots in a p-i-n diode*, *Phys. Rev. B* **68**, 125331 (2003).
- [15] G. Kießlich, A. Wacker, and E. Schöll: *Many-particle charging effects and recombination current through a quantum dot array*, *phys. status solidi (b)* **234**, 215 (2002), excitons and Many-Particle Effects in Semiconductors and Semiconductor Nanostructures (ed. E. Runge and May), in honour of the 60th birthday of R. Zimmermann.
- [16] G. Kießlich, A. Wacker, and E. Schöll: *Sequential tunneling through an array of electrostatically coupled quantum dots*, *Physica E* **12**, 837 (2002).

- [17] G. Kießlich, A. Wacker, and E. Schöll: *Nonlinear transport through an ensemble of quantum dots*, Physica B **314**, 459 (2002).
- [18] A. Rack, R. Wetzler, A. Wacker, and E. Schöll: *Dynamical bistability in quantum-dot structures: Role of Auger processes*, Phys. Rev. B **66**, 165429 (2002).
- [19] R. Wetzler, C. M. A. Kapteyn, R. Heitz, A. Wacker, E. Schöll, and D. Bimberg: *Capacitance-voltage spectroscopy of self-organized InAs/GaAs quantum dots embedded in a pn diode*, phys. status solidi (b) **224**, 79 (2001).
- [20] R. Wetzler, A. Wacker, E. Schöll, C. M. A. Kapteyn, R. Heitz, and D. Bimberg: *Capacitance-voltage characteristics of InAs/GaAs quantum dots embedded in a pn structure*, Appl. Phys. Lett. **77**, 1671 (2000).